


TOP 5 RISKS RELATED TO TAKING THE ABORTION PILL:

1. UTERINE BLEEDING AND HEMORRHAGING
2. INFECTION AND SEPSIS
3. INCREASED RISK OF PELVIC INFLAMMATORY DISEASE
4. MENTAL HEALTH CONDITIONS AND WORSENING PRE-EXISTING CONDITIONS
5. DEATH



Nearly **HALF OF ALL WOMEN** who take the abortion pill seek medical attention after.

Already took the first pill?

REVERSAL MAY STILL BE POSSIBLE.

CALL THE ABORTION PILL REVERSAL HELPLINE:



1-877-558-0333



abortionpillreversal.com

Abortion Pill Reversal* is a safe and effective process that gives your baby a second chance at life.

Thousands of women have successfully reversed the abortion pill and chosen hope over regret.


This 24/7 helpline will connect you with a trusted medical professional nearby who can help you take the next step toward reversing the abortion pill.

Time is critical, and so is your baby.

THIS MESSAGE
BROUGHT
TO YOU BY:

**& THEN NONE
THERE WERE
PROLOVE.COM**

Considering Abortion Pills?



Get the facts. Hear firsthand experiences. Find out what you need to know before you swallow.

**Little
Pills that
Kill**

* IMAGES INSIDE

THE SENSATION OF
A LIFE BEING

Ripped Away


IS AN INDESCRIBABLE HORROR

Hear from former abortion workers.
Hear from patients.

A medication abortion (a.k.a. chemical abortion) uses two drugs to end a pregnancy:

- Mifepristone (first): blocks progesterone, starving the baby.
- Misoprostol (second): causes the uterus to contract and expel the baby.

It is often sold as "a heavy period," but that's not the reality. Listen to real stories from former abortion workers and patients.



"...THEY JUST PUSHED HER OFF
JUST LIKE YOU WERE AT WALMART
OR PURCHASING SOMETHING."

- Bert, former
Planned Parenthood employee



“ I’VE HAD WOMEN
BRING IN A PAD
WITH A
FETUS
LAYING ON IT. ”

- Kylie, former
Planned Parenthood employee

Real photos* of
babies miscarried
or aborted:

*Images have been collected
from clients along with
internet threads/sources



9 Weeks

.75 - 1 INCH



11 Weeks

1.6 - 2 INCHES



13 Weeks

2.7 - 3 INCHES



15 Weeks

~ 6.6 INCHES



17 Weeks

~ 8 INCHES

“ I WAS TOLD TO GO TO
A GROCERY STORE
AND BUY A LEMON TO
SHOW WOMEN HOW
BIG THE BLOOD
CLOTS WOULD GET.”
- S.S., former
Planned Parenthood
employee

